

REPORT DOCUMENTATION PAGE

AFRL-SR-BL-TR-01-

Public reporting burden for this collection of information is estimated to average 1 hour per response, including gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Service, Paperwork Project, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Project, Suite 1204, Arlington, VA 22202-4302.

urces
if this
erson

0480

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE March 2001		3. REPORT TYPE AND DATES COVERED Final report 1 Apr 98 - 31 Mar 01	
4. TITLE AND SUBTITLE (AASERT 98) Student Research in Chemiluminescence				5. FUNDING NUMBERS G-F49620-98-1-0410	
6. AUTHOR(S) Arthur Fontijn					
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Isermann Department of Chemical Engineering Rensselaer Polytechnic Institute 110-8th Street Troy, NY 12180-3590				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING MONITORING AGENCY NAME(S) AND ADDRESS(ES) AFOSR/NA Directorate of Aerospace & Materials Sciences 801 North Randolph Rd., Room 732 Arlington, VA 22203-1977				10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES					
12a. DISTRIBUTION AVAILABILITY STATEMENT Approved for public release distribution is unlimited				12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) Vacuum ultraviolet radiation from rocket plumes offers a unique high-altitude diagnostic observable for identifying and tracking incoming missiles. The interaction of rocket motor combustion products C ₂ H ₂ , C ₂ O, and C ₂ with the atmospheric constituents O and O ₂ was shown to lead to CO(A-X) (4+) chemiluminescence. The spectra and kinetics of the reactions were studied in a heatable pseudo-static photochemical reactor and in fast-flow reactors. The O + C ₂ O and C ₂ + O ₂ reactions were shown to proceed through excited C ₂ O ₂ intermediates. The chemiluminescence from C ₂ + O ₂ was used as the diagnostic for rate coefficient determinations; the results compared well to laser-induced fluorescence measurements, which led to $k(293-1250\text{ K}) = 1.1 \times 10^{-11} \exp(-392\text{ K/T})\text{ cm}^3\text{ molecule}^{-1}\text{s}^{-1}$. In addition to 4+ radiation, the C ₂ + O ₂ reaction produced an apparent continuum in the vuv, probably due to C ₂ O ₂ radiation to a repulsive ground state. This suggests the possibility for developing a chemical laser for the vuv.					
14. SUBJECT TERMS Rocket plumes, CO emission, vacuum ultraviolet, visible radiation, reaction kinetics, C ₂ , O ₂ , C ₂ O, C ₂ H ₂ , O, C ₂ O ₂				15. NUMBER OF PAGES 3	
				16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT UL		

Research Objectives

The purposes of this work were to obtain kinetic information on chemiluminescent reactions for incorporation in missile plume radiation codes and to train students in gas-phase chemiluminescence laboratory techniques.

Results and Publications

The progress made is best described by the publications resulting from the grant.

1. **Arthur Fontijn, Abel Fernandez, Aleksandra Ristanovic, Mai Y. Randall, and Jerome T. Jankowiak, "CO Chemiluminescence and Kinetics of the $C_2 + O_2$ Reaction" *The Journal of Physical Chemistry* 105, xxxx (5 April 2001)**

The reaction $C_2(X^1\Sigma, a^3\Pi) + O_2(X^3\Sigma) \rightarrow CO(X^1\Sigma, A^1\Pi, a'^3\Sigma, d^3\Delta, e^3\Sigma) + CO(X^1\Sigma)$ has been studied in two types of reactors. A Pyrex steady-state fast-flow reactor at approximately 500 K was used to obtain the spectral distributions of the CO ($A^1\Pi-X^1\Sigma$) emissions in the vacuum ultraviolet (vuv) and the CO triplet states emissions in the visible and near ir (vis) wavelength region. The vuv emission had not previously been positively identified. The C_2 was produced from the $C_2Cl_4 + K$ reaction. In a pseudo-static high-temperature photochemistry (HTP) reactor C_2 was made by 193 nm multiphoton dissociation of C_2Cl_4 . That apparatus was used for quenching and rate coefficient experiments in the time domain. The vuv quenching measurements confirm the orbital symmetry argument that the reaction proceeds through excited C_2O_2 intermediates. Reaction schemes for $C_2(X^1\Sigma)$ and for $C_2(a^3\Pi)$ are presented. The results are compared to those from the $O + C_2H_2$ reaction, which leads via $C_2O + O$ to the same band systems emissions, and the differences are discussed below (Paper 2). Measurements of the overall rate coefficients from the decrease of the vis chemiluminescence intensities with time yielded: $k_{vis}(298-711\text{ K}) = 1.1 \times 10^{-11} \exp(-381\text{ K/T}) \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$, with 2σ precision limits of around $\pm 5\%$ and corresponding estimated accuracy limits of about $\pm 21\%$. This expression is in excellent agreement with earlier rate coefficient values determined by different methods, which leads to the recommendation $k(293-1250\text{ K}) = 1.1 \times 10^{-11} \exp(-392\text{ K/T})$. The vuv experiments yielded slightly higher values, the reason for which is uncertain. It is speculated that an apparent continuum observed in the vuv spectra could be 1C_2O_2 excimer radiation to the repulsive ground state. This suggests the possibility of using the $C_2 + O_2$ reaction to produce a chemical laser for vacuum ultraviolet wavelengths.

2. **Arthur Fontijn, Abdellatif Goumri, and Paul E. Brock II, "Pressure-Dependence of the CO($d^3\Delta-a^3\Pi$) Triplet Bands Chemiluminescence Intensities from the O + C₂H₂ Reaction: Mechanistic Implications", *Combustion and Flame*, 121, 699 (2000).**

It is shown that the (visible) Triplet Bands of CO from the O + C₂O reaction decrease in intensity with increasing pressure. This behavior is opposite to that of the (vuv) CO 4+ bands, which increase in intensity. This indicates a mechanism where the triplet emitter molecules form first and by collision-induced cross-relaxation produce the singlet emitters. By contrast the CO 4+ bands from C₂ + O₂ decrease in intensity with increasing pressure and the cross-relaxation mechanism does not contribute significantly there, as further discussed in Paper 1.

Theses

The following theses have resulted in part from support under the present grant:

M.Y. Randall, "Kinetics of the CO Fourth Positive Chemiluminescence from the C₂ + O₂ Reactions", M.Sc., Rensselaer Polytechnic Institute, Troy, NY (1998).

A.S. Blue, "Temperature Dependence of the Kinetics of Transient Species Reactions", Ph.D., Rensselaer Polytechnic Institute, Troy, NY (2000).

Further Students Supported by this Grant

In addition to the two above US students who have completed their theses, one ongoing Ph.D. student, Abel Fernandez, and a dropped-out graduate student Jessica Kasner, participated. A number of Rensselaer US undergraduates also received support under this grant and participated for periods varying from 3 to 12 months. These people are: Jerome T. Jankowiak, Ryan Youngsaye, Todd Alexander, Charles Fung, Christine Brown, Erin Carhart, Inuka Dissanayake, and Michele Nardone. Partially as a result of this first research experience, several of these students decided to go to graduate school.